

An Introduction
to
Applied Partial Differential Equations
Marek Elżanowski

PROBLEMS 2

1. Find the Fourier series representation of:

(a)

$$f(x) = \begin{cases} ax, & -\pi < x \leq 0, \\ bx, & 0 \leq x < \pi. \end{cases} \quad (1)$$

(b)

$$f(x) = e^{ax}, \quad -\pi \leq x \leq \pi. \quad (2)$$

(c)

$$f(x) = \begin{cases} x + 1, & -1 < x < 0, \\ x - 1, & 0 < x < 1. \end{cases} \quad (3)$$

(d)

$$f(x) = \begin{cases} 0, & -2\pi \leq x < -\pi, \\ \sin x, & -\pi \leq x \leq \pi, \\ 0, & \pi \leq x \leq 2\pi. \end{cases} \quad (4)$$

2. Find the Fourier sine series for $f(x) = |\cos x|$ on $0 \leq x \leq \pi$.

3. Find the Fourier series of $f(x) = \pi^2 - x^2$ on $-\pi \leq x \leq \pi$ and use it to find the Fourier series of x and $x(\pi^2 - x^2)$.

4. Find the Fourier sine and cosine series for the function $f(x) = L - x$ on the interval $[0, L]$.

5. Use the Fourier cosine series on the interval $0 \leq x \leq l$ to show that

$$\frac{2}{l} \int_0^l [f(x)]^2 dx = 2a_0^2 + \sum_{n=1}^{\infty} a_n^2. \quad (5)$$

6. (**Bessel's inequality**) Let $f(x)$ be defined on the interval $[-l, l]$, and suppose that $f(x)$ is square integrable on this interval. Assuming that the Fourier series for $f(x)$ exists, show that

$$\frac{a_0^2}{2} + \sum_{n=1}^{\infty} (a_n^2 + b_n^2) \leq \frac{1}{l} \int_{-l}^l |f(x)|^2 dx. \quad (6)$$

7. Use Bessel's inequality for $f(x) = x$ on the interval $[-\pi, \pi]$ to prove that

$$\sum_{n=1}^{\infty} \frac{1}{n^2} \leq \frac{\pi^2}{6}. \quad (7)$$

8. It can be shown that

$$\sum_{n=1}^{\infty} \frac{\sin nx}{\sqrt{n}} \quad (8)$$

converges for all x to an absolutely integrable function on $[-\pi, \pi]$. Use Bessel's inequality to show that there is no function $f(x)$ on $[-\pi, \pi]$ with

$$\int_{-\pi}^{\pi} |f(x)|^2 dx < \infty \quad (9)$$

such that the series (8) is its Fourier series.

9. Assuming that $f(x)$ and $f'(x)$ are defined on $[-l, l]$, show that $f'(x)$ is even if $f(x)$ is odd, and vice versa.

10. Show that if the function $f(x)$ is continuous and piecewise C^1 on the interval $[-l, l]$ with $f(-l) = f(l)$, then the Fourier series of $f'(x)$ can be obtained by differentiating term-by-term the Fourier series of $f(x)$. **Hint:** Integrate by parts the Fourier coefficients of $f'(x)$. Note that the integration by parts will not be valid if $f(x)$ is not continuous. What if we do not require that $f(-l) = f(l)$?

11. Find the complex Fourier series of the following functions:

(a)

$$f(x) = e^x, \quad -\pi < x < \pi, \quad (10)$$

(b)

$$f(x) = \sinh x, \quad -\pi < x < \pi, \quad (11)$$

(c)

$$f(x) = \begin{cases} 0, & 0 < x < 1, \\ 1, & 1 < x < 4. \end{cases} \quad (12)$$

July 10, 2003